

7. Hewitt, J. L. 1913. Twig blight and blossom blight of the apple. *Ark. Agr. Exp. Sta. Bul.* 113:493-505.
8. Lewis, S. and R. N. Goodman. 1965. Mode of penetration and movement of fire blight bacteria in apple leaf and stem tissue. *Phytopath.* 55:719-723.
9. Nightingale, A. A. 1936. Some chemical constituents of apple associated with susceptibility to fire blight. *N. J. Agri. Expt. Sta. Bul.* 613.
10. Schroth, M. N., S. V. Thomson, D. C. Hildebrand, and W. J. Moller. 1974. Epidemiology and control of fire blight. *Ann. Rev. Phytopath.* 12:389-412.
11. Schupp, J. R. and D. C. Ferree. 1987. Deblossoming stimulates second bloom in 'Golden Delicious' apple trees. *Hort. Sci.* 23 (in press).
12. Schupp, J. R. and D. C. Ferree. 1987. The influence of root pruning and deblossoming on the physiology of apple trees. II. photosynthesis, carbohydrates and water rela-
- tions. *J. Amer. Soc. Hort. Sci.* (in press).
13. Shaw, L. 1935. Intercellular humidity in relation to fire blight susceptibility in apple and pear. *Cornell Univ. Agr. Expt. Sta. Memoir* 101.
14. Thompson, J. M. 1972. Fire blight ratings, bloom dates, and precocity of apple varieties tested in the southeast. *Fruit Var. Hort. Digest* 26:84-97.
15. Thomson, S. V., M. N. Schroth, W. J. Moller, and W. O. Reil. 1975. Occurrence of fire blight of pears in relation to weather and epiphytic populations of *Erwinia amylovora*. *Phytopath.* 65:353-358.
16. van der Zwet, T. and H. L. Keil. 1979. Fire Blight: A bacterial disease of *Rosaceous* plants. *USDA Agric. Handbook* 510. U. S. Govt. Printing Off., Washington, D. C. pp. 48-49.
17. van der Zwet, T., W. A. Oitto and H. J. Brooks. 1970. Scoring system for rating the severity of fire blight in pear. *Plant Dis. Repr.* 54:835-839.

Fruit Varieties Journal 42(2):44 1988

Trends of Production, Cultivars and Planting Systems on Apples and Pears in Western Europe

F. WINTER AND M. WELTE

Normal production of 7 million tons from the EEC member countries results in about a 10% over production in a normal crop year in relation to the balance of production to consumption. The top country concerning apple as well as pear production is Italy, whereas in France the new planting rates indicate a reduction. Regarding the area covered by fruit trees within the EEC, since 1977 the apple area decreased from 212,000 ha to less than 200,000 ha and pears also are decreasing. However the small yearly reductions are generally overcompensated by the increase in productivity in modern orchards.

The most important apple cultivars, by country, are as follows: Italy—Golden Delicious 40%, Delicious 20%, and

Imperative 15%; France—Golden Delicious 65%, Delicious 10%, and Granny Smith 8%; F. R. Germany—Golden Delicious, Bos Koop and Cox's Orange; Netherlands—Golden Delicious 30%, Bos Koop and Cox's Orange; Belgium—similar to Netherlands with Jonagold 10%; United Kingdom—Cox's Orange and Bramley's Seedling.

Italy produces more pears than the total of all the other countries with Conference as the main cultivar.

High density planting systems on dwarf to medium rootstocks are more and more preferred in all areas, but individual preference by area do still exist and will continue, influenced by local tradition and ecological reasons—mainly differences in radiation.

From: *Acta Horticulturae* 160:261-68, 1986.