

### Literature Cited

1. Beach, S. A. 1905. The apples of New York. Rpt. of New York Agr. Expt. Sta. 1903. Vol. 1. John Wiley and Son, New York.
2. Fisher, D. V. 1970. Spur strains of McIntosh discovered in British Columbia, Canada. Fruit Var. and Hort Dig. 24:27-32.
3. Heeney, H. B. 1981. New apple cultivars and advanced selections at the Simthfield Experimental Farm. Agriculture Canada, Technical Bull. No. 2, Ministry of Supply and Services, Canada.
4. Lapins, K. O. 1969. Segregation of compact growth types in certain apple seedling progenies. Can. J. Plant Sci. 49:765-768.
5. Lapins, K. O. 1976. Inheritance of compact growth type in apple. J. Amer. Soc. Hort. Sci. 101:133-135.
6. Looney, N. E. and W. D. Lane. 1984. Spur-type growth mutants of McIntosh apple: a review of their genetics, physiology and field performance. Acta Hort. 146:31-46.
7. Lord, W. J., R. A. Damon, Jr. and D. W. Greene. 1983. Variations in growth and productivity among Macspur apple trees, and growth comparisons between spur and nonspur McIntosh and Delicious cultivars. Fruit Var. J. 37:95-99.
8. Lord, W. J., W. J. Bramlage and W. R. Autio. 1985. The Marshall McIntosh apple. Fruit Var. J. 39:37-41.
9. Mehlenbacher, S. A., M. M. Thompson, J. Janick, E. B. Williams, F. H. Emerson, S. S. Korban, D. F. Dayton and L. F. Hough. 1988. 'McShay' apple. HortScience 23:1091-1092.
10. Olien, W. C. 1986. Chick-A-Dee McIntosh: a new spur-type strain. Fruit Var. J. 40:34.
11. Proctor, J. T. A. 1979. Apple cultivars grown in Canada. Fruit Var. J. 33:12-15.
12. Strik, B. C. and J. T. A. Proctor. 1986. Apple cultivars bred in Canada: selections from controlled crosses for commercial production. Fruit Var. J. 40:51-55.
13. Tobutt, K. R. 1984. Breeding columnar apple varieties at East Malling. Scientific Horticulture 35:72-77.
14. Upshall, W. H. 1970. McIntosh, pp. 99-114. In North American apples: varieties, rootstocks, outlook. Michigna State Univ. Press, East Lansing.

## Changing Trends in Blueberry Production and Cultivars in North America

KIM PATTEN<sup>1</sup>

### Introductory Remarks

The blueberry industry in North America has steadily increased over the last few decades. This expansion has occurred throughout North America, encompassing not only the traditional highbush and lowbush blueberry, but also the lowchill highbush, halfhigh, and the rabbiteye blueberry. The latter three types of blueberries represent only a small percentage of the total blueberry production, but have received considerable scientific research aimed at their production and germplasm development.

Concomitant to the rapid expansion of the blueberry industry, has been the development of new germplasm, new cultivars and innovative production practices. To document these

changes the American Society for Horticulture Science and the American Pomological Society sponsored a workshop on this subject in 1989. This workshop chronicled the present status of each type of blueberry, changes in production practices and cultivars, direction for future research and development, and germplasm resources available to meet the future needs of cultivar development. Because of the market demand for the fruit, the expansion of the industry into new production zones, and the continuing endeavors of research to improve blueberry germplasm, the next decade will continue to bring about rapid changes in the North American blueberry industry.

<sup>1</sup>Texas Agricultural Experiment Station, Overton, TX 75684.