

Performance of Trees in the 1990 NC-140 Apple Cultivar/Rootstock Planting: Additional Cultivars and Rootstocks¹

P.M. Hirst, W.R. Autio, J.A. Barden, G.R. Brown, R.M. Crassweller,
P.A. Domoto, and J.R. Schupp.

Abstract

In 1990, up to 10 apple (*Malus X domestica* Borkh.) cultivars were planted on four to seven rootstocks at six sites in the midwestern and eastern United States. The growth and field performance of these trees was measured over 10 years. Although some cultivar x rootstock interactions were evident, tree growth and performance was primarily due to the main effects of cultivar and rootstock. Cultivar had no effect on tree size at three sites, while at other sites 'Rome Beauty', 'Jonagold' and 'McIntosh' trees were the largest. 'Stayman' and 'Empire' trees were among the smallest trees across sites. Trees growing on M.26 EMLA were among the largest trees at all sites, although trees on M.9 EMLA were similar in size at two sites. The smallest trees were produced by B.9, Mark, P.22 and M.27 EMLA rootstocks. No single cultivar produced the highest yield at all sites.

'Golden Delicious' was among the most productive cultivars at three sites, but performed only moderately or poorly at other sites. 'Empire' and 'McIntosh' trees had the lowest yields per tree at most sites. Yields per tree tended to be closely related to tree size, therefore rootstocks producing the largest trees (M.26 EMLA, M.9 EMLA) also produced the largest yields. 'York Imperial' and 'Stayman' trees were highly efficient, and although 'Rome' trees were efficient in some sites, they were inefficient in others. Consistently the most efficient rootstocks were B.9, P.22, and Mark. M.26 EMLA was among the least efficient trees at each site. A significant negative relationship between tree size and yield efficiency was evident at each site, but the relationship differed among sites.

There is a plethora of reports of rootstock performance in the literature, but in most instances, these were conducted with one test cultivar. Studies comparing a number of cultivars growing on a range of rootstocks have produced differing results. Some studies showed that rootstock performance varied depending on the scion cultivar (8, 9) whereas in other studies the rootstock x scion interactions were either insignificant or contributed little to the overall effects (2, 3, 5, 10).

The objective of this study was to compare the growth and performance of a number of apple cultivars growing on up to seven rootstocks across six sites. This paper is one of a series appearing in this issue of the Journal of the American Pomological Society describing results from a

NC-140 cultivar/rootstock trial planted at a number of sites in 1990. Some of the co-operating sites in this planting included cultivars and/or rootstocks additional to those on the main planting (1), and the performance of these additional trees is reported here.

Materials & Methods

An apple planting was established in 1990 as previously described (1, 7). Briefly, the planting consisted of four cultivars ('Smoothee Golden Delicious', 'Nicobel Jonagold', 'Empire', 'Law Rome Beauty') growing on five rootstocks (M.9 EMLA, B.9, Mark, O.3, M.26 EMLA) in 12 sites (CO, IA, IN, KS, KY, MA, ME, OH, PA, TN, UT, VA). The performance of these trees was reported by Autio et al.

¹Purdue University Agricultural Experiment Station Journal paper No. 16507. At participating state agricultural experiment stations, funding was provided by RRF NC-140. We wish to thank the International Dwarf Fruit Tree Association for their financial support of this project and Stark Bro's Nurseries, Louisiana, MO for donating the trees.

Table 1. List of cultivars and rootstocks at each of the six sites. Shaded areas indicate cultivars and rootstocks that formed the main planting described by Autio et al. (1).

| Site | IA | IN | KY | ME | PA | VA |
|-------------------|----|----|----|----|----|----|
| Cultivars | | | | | | |
| Jonathon | X | | | | | |
| Chieftain | X | | | | | |
| Liberty | | X | | | | |
| McIntosh | | | X | X | | |
| York | | | | X | X | |
| Stayman | | | | X | X | |
| Golden Delicious | X | X | X | X | X | X |
| Jonagold | X | X | X | X | X | X |
| Empire | X | X | X | X | X | X |
| Rome | X | X | X | X | X | X |
| Rootstocks | | | | | | |
| P.22 | | X | X | | | |
| M.27 EMLA | | X | | | | |
| M.9 EMLA | X | X | X | | X | X |
| B.9 | X | X | X | X | X | X |
| Mark | X | X | X | X | | X |
| O.3 | X | X | X | X | X | X |
| M.26 EMLA | X | X | X | X | X | X |

(1). Additional trees of a range of cultivars and rootstocks were included in the plantings at six of the sites (Table 1). These were generally combined in a factorial arrangement where each cultivar was growing on each rootstock. For example, in Iowa, the trees additional to the main planting were 'Jonathan' and 'Chieftain' growing on M.9 EMLA, B.9, Mark, O.3, and M.26 EMLA rootstocks.

Each year during October, trunk circumference was measured approximately 25 cm above the graft union, and trunk cross-sectional area was calculated. After defruiting the trees in the first two growing seasons, annual yield per tree was measured. Data collection and analyses were performed by the Massachusetts site cooperator (1). The MIXED procedure of SAS (SAS Institute, Cary, NC) was used to analyze the data. For a more complete description of statistical procedures, see Autio et al. (1).

Results & Discussion

Tree size. Tree size was the product of cultivar, rootstock, and their interaction, although this was not consistent across all sites. At three sites (IN, KY, ME) cultivar had no effect on tree size while the interaction between cultivar and rootstock was significant at some sites (IA, IN, ME, PA) but not at others (KY, VA) (Tables 2-7). 'Jonagold', 'Rome' and 'Golden Delicious' consistently ranked among the largest trees at each site, while 'Empire' and 'Stayman' tended to be the smallest. As expected, trees on M.26 EMLA rootstock ranked as the largest at each site, although at two sites, those on M.9 EMLA and O.3 were similar in size. Trees growing on Mark, B.9, P.22, and M.27 EMLA were similar in size and were consistently in the smallest category at every site. At all sites, trees on B.9 were smaller than those on M.9 EMLA.

Yield per tree. In ME, cultivar had no effect on yield per tree, but cultivar differences were evident at all other sites. There was no cultivar that produced high yields at all sites. 'Golden Delicious' was among the most productive in three sites (IA, IN, KY) but at other sites, was intermediate




Figure 1. Relationships between trunk cross-sectional area and cumulative yield efficiency (1992-99) of various apple cultivars growing on a range of rootstocks in six sites. The relationships were: IA $y=2.79-0.017x$, $r^2=0.85^{*}$; IN, $y=2.076-0.006x$, $r^2=0.21^{*}$; KY, $y=3.44-0.012x$, $r^2=0.52^{***}$; ME, $y=4.15-0.022x$, $r^2=0.30^{*}$; PA, $y=3.89-0.019x$, $r^2=0.15^{*}$; VA, $y=5.19-0.015x$, $r^2=0.56^{***}$.**

Table 2. Trunk cross-sectional area and yield as affected by cultivar and rootstock after 10 years in the 1990 NC-140 Cultivar/Rootstock Trial in Iowa. All values are least-squares means, adjusted for missing subclasses. Cultivar and rootstock interacted significantly to affect trunk cross-sectional area, yield per tree, and yield efficiency so mean separations are presented for rootstock within each cultivar.²

| Rootstock | Golden Delicious | Jonagold | Empire | Rome | Jonathon | Chieftain | Mean |
|---|------------------|----------|---------|---------|----------|-----------|--------|
| <i>Trunk cross-sectional area (cm²)</i> | | | | | | | |
| M.9 EMLA | 89 b | 105 ab | 65 b | 86 b | 94 a | 81 b | 87 b |
| B.9 | 48 c | 51 c | 35 c | 58 c | 49 b | 44 c | 47 c |
| Mark | 40 c | 44 c | 29 c | 50 c | 36 b | 44 c | 41 c |
| O.3 | 95 ab | 88 b | 83 ab | 101 ab | 94 a | 78 b | 90 b |
| M.26 EMLA | 116 a | 124 a | 94 a | 114 a | 109 a | 121 a | 113 a |
| Mean | 78 ab | 82 a | 61 b | 82 a | 76 ab | 74 ab | |
| <i>Cumulative yield per tree (1992-99, kg)</i> | | | | | | | |
| M.9 EMLA | 122 ab | 99 a | 121 a | 107 a | 102 ab | 104 a | 109 a |
| B.9 | 105 bc | 83 ab | 74 b | 103 a | 89 b | 89 a | 90 bc |
| Mark | 84 c | 57 b | 70 b | 89 a | 80 b | 96 a | 79 c |
| O.3 | 137 a | 85 ab | 131 a | 104 a | 122 a | 108 a | 115 a |
| M.26 EMLA | 96 bc | 78 ab | 93 b | 99 a | 90 b | 105 a | 94 b |
| Mean | 109 a | 81 b | 98 ab | 100 a | 97 ab | 100 a | |
| <i>Cumulative yield efficiency (1992-99, kg/cm² TCA)</i> | | | | | | | |
| M.9 EMLA | 1.39 a | 0.95 bc | 1.91 bc | 1.27 bc | 1.14 b | 1.29 bc | 1.33 b |
| B.9 | 2.28 a | 1.67 a | 2.22 ab | 1.80 ab | 1.91 a | 2.11 a | 2.00 a |
| Mark | 2.08 a | 1.29 ab | 2.58 a | 1.87 a | 2.29 a | 2.25 a | 2.06 a |
| O.3 | 1.46 b | 1.07 bc | 1.61 c | 1.05 c | 1.32 b | 1.39 ab | 1.32 b |
| M.26 EMLA | 0.83 c | 0.65 c | 1.02 d | 0.88 c | 0.84 b | 0.89 c | 0.85 c |
| Mean | 1.61 ab | 1.12 c | 1.87 a | 1.38 bc | 1.50 bc | 1.59 b | |

²Separation among overall rootstock means and among overall cultivar means by Tukey's HSD ($P = 0.05$). Mean separation among rootstocks within cultivars by t test ($P = 0.05$) with a Bonferroni adjustment (adjusted $P = 0.005$).

(PA) or low (VA) in terms of yield per tree of other cultivars. 'Rome' was relatively productive in IA, IN, PA, and VA but was among the lowest yielding cultivars in KY. 'York' was also quite productive at the two sites where this cultivar was included (PA and VA). Trees growing on M.26 EMLA and M.9 EMLA rootstocks tended to be the highest yielding at each site, except in Iowa where trees on M.26 EMLA were relatively unproductive. The smallest trees also tended to have the lowest yields per tree, therefore typically trees growing on

Mark, B.9, P.22, and M.27 EMLA rootstocks had the lowest yields at each site.

Yield efficiency. There were no cultivars that had consistently high yield efficiencies across all sites. For example, 'Rome' was among the most efficient cultivars in IN, ME, PA, and VA but was among the least efficient in IA and KY. 'York' and 'Stayman' trees were highly efficient while 'McIntosh' trees had low efficiency, although these cultivars were only planted at two sites. Overall, trees on B.9 were the most efficient, although trees

Table 3. Trunk cross-sectional area and yield as affected by cultivar and rootstock after 10 years in the 1990 NVC-140 Cultivar/Rootstock Trial in Kentucky. All values are least-squares means, adjusted for missing sub-classes. Cultivar and rootstock did not interact significantly to affect trunk cross-sectional area, yield per tree, or yield efficiency, so separations are presented only for overall rootstock and cultivar means.²

| Rootstock | Golden Delicious | Jonagold | Empire | Rome | Liberty | Mean |
|---|------------------|----------|---------|--------|---------|---------|
| <i>Trunk cross-sectional area (cm²)</i> | | | | | | |
| M.9 EMLA | — | — | 63 | 104 | 112 | 99 a |
| B.9 | 61 | — | 37 | 63 | 60 | 56 b |
| Mark | 32 | 20 | 29 | 35 | 35 | 35 b |
| O.3 | — | 103 | 46 | — | — | 86 ab |
| M.26 EMLA | 176 | 162 | 108 | 128 | 118 | 132 a |
| P.22 | — | 42 | 19 | 36 | 24 | 33 b |
| Mean | 85 a | 82 a | 54 a | 74 a | 74 a | |
| <i>Cumulative yield per tree (1992-99, kg)</i> | | | | | | |
| M.9 EMLA | — | — | 142 | 122 | 268 | 197 a |
| B.9 | 168 | — | 104 | 128 | 158 | 139 a |
| Mark | 96 | 76 | 83 | 67 | 98 | 90 b |
| O.3 | — | 181 | 131 | — | — | 173 ab |
| M.26 EMLA | 252 | 220 | 182 | 179 | 269 | 219 a |
| P.22 | — | 116 | 60 | 78 | 93 | 86 b |
| Mean | 175 ab | 154 ab | 119 b | 122 b | 184 a | |
| <i>Cumulative yield efficiency (1992-99, kg/cm² TCA)</i> | | | | | | |
| M.9 EMLA | — | — | 3.17 | 1.25 | 2.52 | 2.21 ab |
| B.9 | 2.72 | — | 3.59 | 2.26 | 3.69 | 3.04 a |
| Mark | 2.75 | 3.10 | 3.13 | 1.92 | 3.74 | 2.89 a |
| O.3 | — | 1.78 | 2.65 | — | — | 1.99 ab |
| M.26 EMLA | 1.57 | 1.55 | 1.94 | 1.36 | 2.43 | 1.76 b |
| P.22 | — | 2.72 | 3.28 | 2.22 | 3.90 | 2.99 a |
| Mean | 2.22 ab | 2.38 ab | 2.90 ab | 1.75 b | 3.13 a | |

²Separation among overall rootstock means and among overall cultivar means by Tukey's HSD ($P = 0.05$).

growing on P.22 rootstock were similar at the two sites where this rootstock was included. Trees growing on Mark were also consistently efficient, as opposed to M.26 EMLA, which was the least efficient rootstock at each site.

At each site, there was a significant negative relationship between tree size and yield efficiency (Figure 1). The slopes defining these relationships were broadly similar across sites. Trees in VA were gen-

erally much more efficient than those at other sites, due to higher yields from trees of similar sizes. Strong relationships between tree size and yield efficiency have previously been described where smaller trees were more efficient (3, 4, 6).

When trees were grouped by cultivar (across all sites and rootstocks) or by rootstock (across all sites and cultivars), there were not significant relationships between cumulative yield efficiency and tree size in

Table 4. Trunk cross-sectional area and yield as affected by cultivar and rootstock after 10 years in the 1990 NVC-140 Cultivar/Rootstock Trial in Maine. All values are least-squares means, adjusted for missing sub-classes. Cultivar and rootstock interacted significantly to affect trunk cross-sectional area, yield per tree, or yield efficiency, so mean separations are presented for rootstock within each cultivar.^z

| Rootstock | Golden Delicious | Jonagold | Empire | Rome | McIntosh | Mean |
|---|------------------|----------|--------|---------|----------|---------|
| <i>Trunk cross-sectional area (cm²)</i> | | | | | | |
| B.9 | — | 29 b | 33 bc | 28 | 35 b | 32 c |
| Mark | 40 b | 63 a | 31 c | 47 a | 42 ab | 44 b |
| O.3 | 61 ab | — | 57 ab | 46 a | — | 58 ab |
| M.26 EMLA | 69 a | 65 a | 61 a | 41 a | 71 a | 61 a |
| Mean | 51 a | 54 a | 45 a | 40 a | 52 a | |
| <i>Cumulative yield per tree (1992-99, kg)</i> | | | | | | |
| B.9 | — | 95 b | 99 b | 123 a | 103 a | 103 b |
| Mark | 138 a | 209 a | 102 b | 160 a | 89 a | 138 a |
| O.3 | 191 a | — | 184 a | 160 a | — | 182 a |
| M.26 EMLA | 165 a | 184 a | 151 a | 133 a | 129 a | 152 a |
| Mean | 152 a | 173 a | 134 a | 146 a | 114 a | |
| <i>Cumulative yield efficiency (1992-99, kg/cm² TCA)</i> | | | | | | |
| B.9 | — | 3.32 a | 3.30 a | 4.43 a | 2.75 a | 3.44 a |
| Mark | 3.53 a | 3.33 a | 3.18 a | 3.49 ab | 2.25 a | 3.15 ab |
| O.3 | 3.13 ab | — | 3.36 a | 3.48 ab | — | 3.14 ab |
| M.26 EMLA | 2.39 b | 2.98 a | 2.63 a | 3.24 b | 1.85 a | 2.63 b |
| Mean | 3.13 a | 3.22 a | 3.12 a | 3.71 a | 2.28 b | |

^zSeparation among overall rootstock means and among overall cultivar means by Tukey's HSD ($P = 0.05$). Mean separation among rootstocks within cultivars by t test ($P=0.05$) with a Bonferroni adjustment (adjusted $P = 0.008$).

most cases (data not presented). This suggests that site was the predominant influence on tree performance in this study. Site also had a much larger effect than rootstock on tree performance in a recent study with 'Gala' growing on 18 dwarf and 4 semi-dwarf rootstocks (6). This reiterates the importance of conducting coordinated trials such as this to enable the response of cultivars and/or rootstocks to be measured across widely varying sites. Such trials are a necessary pre-requisite to making appropriate site-specific recommendations to growers.

Literature cited

1. Autio, W. R., J. L. Anderson, J. A. Barden, G. R. Brown, R. M. Crassweller, P. A. Domoto, A. Erb, D. C. Ferree, A. Gaus, P. M. Hirst, C. A. Mullins and J. R. Schupp 2001. Performance of 'Golden Delicious', 'Jonagold', 'Empire', and 'Rome' apple trees on five rootstocks over ten years in the 1990 NC-140 cultivar/rootstock trial. *J. Amer. Pom. Soc.* 55:xx-xx
2. Barritt, B.H., B.S. Konishi, and M.A. Dilley. 1995. Performance of three apple cultivars with 23 dwarfing rootstocks during 8 seasons in Washington. *Fruit Var. J.* 49:158-170.
3. Ferree, D. C., P. M. Hirst, J. C. Schmid and P. E. Dotson 1995. Performance of three apple

Table 5. Trunk cross-sectional area and yield as affected by cultivar and rootstock after 10 years in the 1990 NC-140 Cultivar/Rootstock Trial in Pennsylvania. All values are least-squares means, adjusted for missing subclasses. Cultivar and rootstock interacted significantly to affect trunk cross-sectional area, yield per tree, and yield efficiency so mean separations are presented for rootstock within each cultivar.²

| Rootstock | Golden Delicious | Jonagold | Empire | Rome | McIntosh | York | Stayman | Mean |
|---|------------------|----------|--------|--------|----------|---------|---------|--------|
| <i>Trunk cross-sectional area (cm²)</i> | | | | | | | | |
| M.9 EMLA | 70 b | 86 a | 52 b | 72 b | 90 a | 62 bc | 41 b | 67 b |
| B.9 | 45 c | 51 b | 39 b | 52 b | 51 b | 45 c | 40 b | 46 c |
| O.3 | 68 b | 62 b | 57 b | 69 b | 45 b | 78 ab | 60 b | 63 b |
| M.26 EMLA | 96 a | 97 a | 88 a | 119 a | 110 a | 85 a | 85 a | 97 a |
| Mean | 70 ab | 74 ab | 59 b | 78 a | 74 ab | 67 ab | 56 b | |
| <i>Cumulative yield per tree (1992-99, kg)</i> | | | | | | | | |
| M.9 EMLA | 149 a | 124 a | 102 a | 327 ab | 131 a | 206 a | 122 b | 166 a |
| B.9 | 121 a | 103 a | 77 a | 265 c | 113 ab | 179 a | 134 b | 142 b |
| O.3 | 162 a | 114 a | 127 a | 311 bc | 77 b | 203 a | 201 a | 171 a |
| M.26 EMLA | 147 a | 115 a | 127 a | 268 a | 112 ab | 184 a | 169 ab | 175 a |
| Mean | 145 cd | 114 de | 108 e | 318 a | 108 e | 193 b | 156 c | |
| <i>Cumulative yield efficiency (1992-99, kg/cm² TCA)</i> | | | | | | | | |
| M.9 EMLA | 2.20 ab | 1.50 a | 1.98 a | 4.57 a | 1.52 b | 3.37 b | 3.00 a | 2.59 b |
| B.9 | 2.74 a | 2.09 a | 1.99 a | 5.16 a | 2.58 a | 4.42 a | 3.58 a | 3.22 a |
| O.3 | 2.42 a | 1.90 a | 2.22 a | 4.53 a | 1.81 ab | 2.69 bc | 3.43 a | 2.71 b |
| M.26 EMLA | 1.57 b | 1.24 a | 1.47 a | 3.13 b | 1.07 b | 2.19 c | 2.05 b | 1.82 c |
| Mean | 2.24 bc | 1.68 c | 1.91 c | 4.35 a | 1.74 c | 3.17 b | 3.01 b | |

²Separation among overall rootstock means and among overall cultivar means by Tukey's HSD ($P = 0.05$). Mean separation among rootstocks within cultivars by t test ($P = 0.05$) with a Bonferroni adjustment (adjusted $P = 0.008$).

Table 6. Trunk cross-sectional area and yield as affected by cultivar and rootstock after 10 years in the 1990 NC-140 Cultivar/Rootstock Trial in Virginia. All values are least-squares means adjusted for missing sub-classes. Cultivar and rootstock did not interact significantly to affect trunk cross-sectional area, yield per tree, or yield efficiency, so separations are presented only for overall rootstock and cultivar means.²

| Rootstock | Golden Delicious | Jonagold | Empire | Rome | York | Stayman | Mean |
|---|------------------|----------|--------|---------|---------|---------|---------|
| <i>Trunk cross-sectional area (cm²)</i> | | | | | | | |
| M.9 EMLA | 109 | 125 | 96 | 126 | 106 | 112 | 112 b |
| B.9 | 64 | 65 | 52 | 63 | 57 | 45 | 58 c |
| Mark | 58 | 69 | 65 | 86 | 81 | 44 | 67 c |
| O.3 | 104 | 132 | 125 | 110 | 114 | 92 | 113 b |
| M.26 EMLA | 117 | 153 | 155 | 153 | 161 | 144 | 147 a |
| Mean | 90 ab | 109 a | 99 abc | 108 ab | 104 abc | 88 b | |
| <i>Cumulative yield per tree (1992-99, kg)</i> | | | | | | | |
| M.9 EMLA | 348 | 417 | 331 | 498 | 396 | 472 | 410 ab |
| B.9 | 241 | 309 | 218 | 309 | 273 | 238 | 264 c |
| Mark | 200 | 262 | 201 | 310 | 247 | 202 | 237 c |
| O.3 | 314 | 353 | 343 | 412 | 431 | 390 | 374 b |
| M.26 EMLA | 355 | 456 | 371 | 553 | 439 | 439 | 435 a |
| Mean | 291 c | 360 ab | 293 c | 416 a | 357 ab | 348 bc | |
| <i>Cumulative yield efficiency (1992-99, kg/cm² TCA)</i> | | | | | | | |
| M.9 EMLA | 3.23 | 3.47 | 3.84 | 3.94 | 3.84 | 4.26 | 3.76 bc |
| B.9 | 3.82 | 4.79 | 4.22 | 4.88 | 4.87 | 5.24 | 4.64 a |
| Mark | 3.48 | 4.03 | 3.91 | 3.66 | 3.23 | 4.52 | 3.81 b |
| O.3 | 2.99 | 2.62 | 2.86 | 3.65 | 3.78 | 4.21 | 3.35 cd |
| M.26 EMLA | 3.09 | 2.99 | 2.47 | 3.67 | 2.83 | 3.08 | 3.02 d |
| Mean | 3.32 b | 3.58 ab | 3.46 b | 3.96 ab | 3.71 ab | 4.26 a | |

²Separation among overall rootstock means and overall cultivar means by Tukey's HSD ($P = 0.05$).

cultivars with 22 dwarfing rootstocks during 8 seasons in Ohio. *Fruit Var. J.* 49:171-178.

4. Hirst, P. M. and D. C. Ferree 1995. Rootstock effects on shoot morphology and spur quality of 'Delicious' apple and relationships with precocity and productivity. *J. Amer. Soc. Hort. Sci.* 120:622-634.
5. Hirst, P. M. and D. C. Ferree 1995. Effect of rootstock and cultivar on the growth and precocity of young apple trees. *Fruit Var. J.* 49:34-41.
6. Hirst, P. M. and NC-140 cooperators. 2001. Early performance of 'Gala' apple on 18 dwarf and 4 semi-dwarf rootstocks growing at 24 sites in North America. *Acta Hort.* (in press).
7. NC-140. 1996. Rootstock and scion cultivar interact to affect apple tree performance: A

five-year summary of the 1990 NC-140 cultivar/rootstock trial. *Fruit Var. J.* 50:175-187.

8. Schupp, J. R. 1995. Growth and performance of four apple cultivars on M.26 and Mark rootstocks, with or without preplant mineral nutrients. *Fruit Var. J.* 49:198-204.
9. Schupp, J. R. and S. I. Koller 1997. Growth and productivity of four summer ripening disease-resistant apple cultivars on M.27 EMLA, M.26 EMLA and Mark rootstocks. *Fruit Var. J.* 51:161-164.
10. Schupp, J. R. and S. I. Koller 1998. Growth and productivity of disease-resistant apple cultivars on M.27 EMLA, M.26 EMLA and Mark rootstocks. *Fruit Var. J.* 52:140-154.