

# Evaluation of a Spray-on Product for Improving Red Color and Decreasing Sunburn Damage of 'Red Chief Delicious' Apples

T. SOTIROPOULOS<sup>1</sup>, N. VOULGARAKIS<sup>2</sup>, D. TRIANTAFILLOU<sup>2</sup>, I. MANTHOS<sup>3</sup>, P. XAFAKOS<sup>4</sup>

**Additional index words.** Fruit quality, Firmness, Fruit color, Physiological disorders, Solar radiation.

## Abstract

Sunburn is a severe type of solar radiation damage that causes meaningful economic losses in several vegetables and fruits, including apples. In this study, we examined the efficacy of a new product, Parasol, for protecting apples (cv. 'Red Chief Delicious') against sunburn. Specifically, we examined the effect of Parasol spraying on fruit quality attributes, including fruit color, flesh firmness, soluble solids and acidity at harvest and after a 5-month period of storage. Spraying apple trees with Parasol resulted in better red fruit coloration and reduced sunburn symptoms. Parasol (2.5 and 5 L/t) treatments resulted in an increase of fruit flesh firmness compared to the control, for both years. Fruit weight, soluble solids concentration and acidity did not differ significantly between treatments. Hue values showed that apples from the control treatment had significantly less red color than the Parasol treatments. The intensity of the red color decreased in the following order: Parasol (5 L/t) > Parasol (2.5 L/t) > Control. From our results it is concluded that Parasol protects apples from sunburn injury, increases red coloration and fruit firmness and therefore their commercial value.

## Introduction

Apple (*Malus × domestica* Borkh.) is one of the most important fruit crops cultivated in temperate zones of the world. Among fruit quality attributes, apple skin color is a crucial component that determines apple marketability and any factors that adversely influence the appearance decreases the value of the fruits. Accordingly, apple sunburn, a fruit discoloration caused by excessive solar radiation and heat, is a common defect decreasing the value of affected fruits in hot dry weather conditions and reduces post-harvest life (Felicetti and Schrader, 2009; Schrader *et al.* 2003). Losses in apple production caused by sunburn in Washington State (U.S.A.) average about 10% if no preserving measures are taken (Schmidt, 2018). In Australia, losses may range from

40 to 50%, particularly for delicate apple cultivars (Lolicato, 2011) and in Chile 13%. 'Granny Smith' and 'Gala' are considered to be most susceptible to sun damage. Other susceptible apple cultivars are 'Jonagold', 'Braeburn', 'Golden Supreme', 'Ginger Gold' and 'Fuji' (Evans, 2004). According to Schrader *et al.* (2003) 'Pink Lady' is more tolerant than 'Cameo' and 'Honeycrisp'. However, the severity of the symptoms differs from year to year depending on climatic conditions.

Apple cultivars grown mainly in lowland areas are particularly vulnerable to sunburn. The apple cultivar 'Red Chief Delicious', when cultivated in low-lying regions in Greece, is sensitive to sunburn. The injury on the fruit skin becomes visible either as a stain, or in more intense cases as a brown

<sup>1</sup> Hellenic Agricultural Organization 'Demeter', Institute of Plant Breeding and Genetic Resources, Department of Deciduous Fruit Tree Growing in Naoussa, 59035 Naoussa, Greece.

<sup>2</sup> International University of Greece, Department of Supply Chain Management, 60100 Katerini, Greece.

<sup>3</sup> Hellenic Agricultural Organization 'Demeter', Institute of Plant Breeding and Genetic Resources, Department of Nut Trees, 35100 Neo Krikello-Lamia, Greece

<sup>4</sup> Skydra Pellas, 58500 Greece

<sup>5</sup> Corresponding author: thosotir@otenet.gr

necrotic blotch, leading ultimately to entire loss of the commercial value of the fruit.

The categories of sunburn are classified as:

i) Sunburn necrosis appears as a depressed dark brown or necrotic blotch on the part of fruit exposed to the sun. It is generally caused by heat induced in fruit by direct sunlight, when the fruit skin temperature of an apple reaches  $52^{\circ}\pm 1^{\circ}\text{C}$  for  $\geq 10$  minutes, ii) Sunburn browning appears as a yellow or brown blotch on the side of the fruit exposed to the sun. Cells do not die and damage first appears externally, although deeper layers may show further damage after cool storage. Sunburn browning in apples usually starts to appear when fruit skin temperatures rises up to  $46^{\circ}$  to  $49^{\circ}\text{C}$  for one hour, depending on various cultivars, iii) Photo-oxidative sunburn. Initially it appears as white, faded skin in a patch on the side of fruits exposed to the sun, that eventually becomes brown and occasionally also necrotic. It can take place at more or less low air temperature and already at a skin temperature of less than  $45^{\circ}\text{C}$  (Fellicetti and Schrader, 2008; Racsko and Schrader, 2012).

Delayed sunburn symptoms are similar to sunburn browning and can broaden or worsen within the first four months of fruits in cold storage. Fruit that had pre-harvest sunburn that later covered over during red color development are not suitable for long-term storage (Racsko and Schrader, 2012). Beyond sunburn symptoms previously mentioned, heat and light stress can cause other fruit imperfections and disorders. External disorders can be detected such as cracking, crinkle, 'Fuji' stain, hot summer stress disorder, lenticel marking, russetting and splitting. Furthermore, internal disorders occur such as water core, bitter bit and heat stress internal browning.

There are several actions that can be taken to protect apples against sunburn. These may involve the use of climate ameliorating techniques such as the use of cultivars that are more tolerant to sunburn, the scheduling of irrigations to avoid tree water stress, the

training of trees to develop an appropriate canopy, the avoidance of excessive summer pruning, the use of cover cropping and the improvement of air movement through the fruit block. Other measures involve the use of shade nettings, over-tree sprinkler cooling systems and spray-on products such as kaolin, calcium carbonate, wax and talc-based products. Ascorbic acid, abscisic acid and anti-transpirant compounds were tested as well (Amarante *et al.*, 2011; Evans and van der Gulik, 2011; Glenn 2009; Iams *et al.*, 2009; Lal and Sahu, 2017, Lolicato, 2011, Reig *et al.*, 2016; Schupp *et al.*, 2002). The objective of this research was to examine the efficacy of two rates of the spray-on product Parasol in protecting apple fruit from sunburn. In addition, we tested whether Parasol affects the quality attributes of the fruit. The apple cultivar 'Red Chief Delicious' was chosen for this research, since it is commonly accepted among consumers in Greece.

### Materials and Methods

Trees of the apple cv. 'Red Chief Delicious' grafted onto M.26 rootstock were selected for this research. The trees were 13 years old, planted in a randomized complete block design with  $3.5 \times 2.5$  m spacing and trained to a palmette system. Twenty-five trees were used for each treatment (five replications x five-tree units). The trees received standard horticultural practices regarding pruning, irrigation and fertilization.

Apple trees were sprayed by an airblast sprayer with 2.5L and 5L Parasol (Nature S.A. Nea Efessos, Pieria, Greece) per 1000L of water for two consecutive years (2017 and 2018) in an apple orchard located in Naoussa (northern Greece, long.  $22^{\circ}12'0''$  E; lat.  $40^{\circ}29'04''$  N; elevation 350 m) on the following dates: 15 July, 28 July, and 19 Aug. (one month prior to harvest). Control trees were not sprayed. Parasol is a dense, liquid calcium carbonate dispersion product, based on natural, ultrafine particles of  $1.4 \mu\text{m}$ . When Parasol is applied to plants, a very



**Table 2.** The influence of two rates of Parasol on mean fruit weight, soluble solids concentration, acidity, and firmness of 'Red Chief Delicious' fruit at harvest in 2017 and 2018.

Year	Treatment	Fruit weight (g)	Soluble solids (%)	Acidity (% malic acid)	Fruit Firmness (N)
2017	Control	284.9 ± 35.9 a <sup>z</sup>	13.88 ± 3.2 a	0.23 ± 0.01 a	64.14 ± 14.71 b
	Parasol (2.5l/t)	283.7 ± 31.7 a	13.18 ± 2.8 a	0.25 ± 0.02 a	67.86 ± 17.65 a
	Parasol (5l/t)	289.4 ± 32.4 a	13.70 ± 2.6 a	0.24 ± 0.02 a	69.14 ± 13.73 a
2018	Control	265.3 ± 24.9 a	12.70 ± 2.3 a	0.18 ± 0.02 a	61.98 ± 18.63 b
	Parasol (2.5l/t)	267.3 ± 30.4 a	12.30 ± 2.9 a	0.20 ± 0.03 a	69.04 ± 14.70 a
	Parasol (5l/t)	264.9 ± 31.5 a	11.20 ± 2.0 b	0.21 ± 0.04 a	69.73 ± 8.82 a

<sup>z</sup> Means (± SD) within columns and years followed by common letters are not significantly different (Tukey's test,  $P \leq 0.05$ )

dates: 10 June (1000), 14 June (1000), 16 June (1024), 24 June (1042) and 19 July (1022). Maximum temperatures in 2017, were recorded on the following dates: 27 June (38°C), 1 July (38°C), 2 July (39°C), 12 July (38°C), 3 Aug. (38°C), 6 Aug. (38°C). In 2018, maximum temperatures were recorded on the following dates: 5 June (33°C), 13 July (35°C), 21 July (35.5°C), 30 July (35°C), 7 Aug. (35.5°C), 8 Aug. (35.5°C) and 19 Aug. (36°C). Lastly, the 2018 rainfall was higher than in 2017.

In 2017, Parasol at (2.5 and 5 L/t) increased fruit flesh firmness compared to the control (Table 2). Soluble solids concentration, mean fruit weight, and acidity did not differ

significantly among treatments. However, in 2018, soluble solids concentration was lower for fruit from the Parasol (5 L/t) treatment compared to the control. To examine whether application of Parasol on apple trees affects fruit coloration in comparison to the control, the color of apples was measured. Lightness (L) and chroma (C) did not differ significantly between treatments (Table 3). However, Hue values showed that control fruit had significantly less red color than the Parasol treatments. The intensity of the red color decreased in the following order: Parasol (5 L/t) > Parasol (2.5 L/t) > Control (Table 3). The red color development of apples increased for trees treated with Parasol due

**Table 3.** The influence of two rates of Parasol on color parameters L, a, b, Chroma (C) and Hue angle of 'Red Chief Delicious' fruit at harvest in 2017 and 2018.

Year	Treatment	L	a	b	C	Hue
2017	Control	41.22 ± 3.2 az	25.01 ± 2.6 c	21.54 ± 2.5 a	33.01 ± 4.1 a	40.69 ± 4.8 a
	Parasol (2.5l/t)	43.72 ± 3.3 a	27.80 ± 3.2 b	18.82 ± 2.1 b	33.57 ± 4.5 a	34.21 ± 4.0 b
	Parasol (5l/t)	43.95 ± 4.0 a	30.40 ± 3.6 a	16.09 ± 1.9 c	34.39 ± 4.0 a	27.92 ± 3.6 c
2018	Control	40.25 ± 2.8 a	28.93 ± 2.8 b	26.36 ± 2.6 a	39.14 ± 5.2 a	42.30 ± 5.1 a
	Parasol (2.5l/t)	39.57 ± 2.3 a	34.21 ± 4.1 a	19.44 ± 2.4 b	39.35 ± 4.6 a	29.68 ± 3.6 b
	Parasol (5l/t)	42.49 ± 3.0 a	35.32 ± 2.9 a	17.75 ± 2.0 c	39.53 ± 4.4 a	26.56 ± 3.9 c

<sup>z</sup> Means (± SD) within columns and years followed by common letters are not significantly different (Tukey's test,  $P \leq 0.05$ ).

**Table 4.** The influence of two rates of Parasol on soluble solids concentration, acidity, and firmness of ‘Red Chief Delicious’ apple fruit after a five-months of cold storage in two years.

Year	Treatment	Soluble solids (%)	Acidity (% malic acid)	Fruit firmness (N)
2017	Control	14.81 ± 2.1 a <sup>z</sup>	0.20 ± 0.03 a	50.41 ± 7.85 b
	Parasol (2.5l/t)	14.38 ± 1.8 a	0.22 ± 0.02 a	57.47 ± 9.81 a
	Parasol (5l/t)	14.59 ± 1.6 a	0.20 ± 0.03 a	58.74 ± 8.83 a
2018	Control	13.77 ± 1.6 a	0.15 ± 0.02 a	51.39 ± 5.88 b
	Parasol (2.5l/t)	13.49 ± 2.1 a	0.17 ± 0.03 a	60.70 ± 4.90 a
	Parasol (5l/t)	12.27 ± 1.4 b	0.16 ± 0.02 a	60.21 ± 5.88 a

<sup>z</sup>Means (± SD) within columns and years followed by common letters are not significantly different (Tukey’s test,  $P \leq 0.05$ ).

to the lower discoloration percentage as was shown in Table 5 and protection of high solar radiation on the fruit’s skin.

We further examined fruit quality attributes of apples stored for 5 months in cooling chambers. The fruit soluble solids concentration increased with respect to the harvest stage (Table 4). Treatments with Parasol (2.5 and 5 L/t) had increased fruit firmness in comparison to the control.

During the 2nd year, the fruit soluble solids concentration increased slightly during storage (Table 4), and treatment differences with regard to the fruit firmness, and soluble solids concentration and acids at harvest were maintained during cold storage for 5 months. At harvest in both years, more from the

control treatment had sunburns and light discoloration compared to both Parasol treatments (Table 5). Parasol treatment (5 L/t) had the most fruit in Class I (Fig. 1). Alvarez *et al.* (2015) reported that calcium carbonate has a thermoprotector effect on apples and that the degree of protection depends on the reflectance and the amount of residue settled on the surface. Glenn *et al.* (2002) also reported that ‘Crimson seedless’ grapevines treated with  $\text{CaCO}_3$  were less prone to sunburn damage than untreated grapes due to reduced fruit temperature and exposure to UV radiation because  $\text{CaCO}_3$  reflected UV radiation. Furthermore, Ahmed *et al.* (2011) reported that grapevines treated with  $\text{CaCO}_3$  were less prone to sunburn damage than

**Table 5.** The influence of two rates of Parasol on the percentage of ‘Red Chief Delicious’ apples assigned to three classifications based on the color and sunburn injury at harvest in two years.

Year	Treatment	Class I*	Class II	Class III
2017	Control	68 c <sup>z</sup>	19 a	13 a
	Parasol (2.5l/t)	86 b	10 b	4 b
	Parasol (5l/t)	92 a	6 c	2 c
2018	Control	72 c	16 a	12 a
	Parasol (2.5l/t)	87 b	8 b	5 b
	Parasol (5l/t)	93 a	4 c	3 c

<sup>z</sup>Means (± SD) within columns and years followed by common letters are not significantly different (Tukey’s test,  $P \leq 0.05$ ).

\* Class I (extra quality-absence of any sunburn symptoms), Class II (light discoloration) and Class III (visual sunburn spots).



**Figure 1.** ‘Red Chief Delicious’ fruit coloration at harvest as affected by two rates of Parasol applied foliarly three times from 8 to 4 weeks before harvest.

untreated grapes. Previous researchers found that  $\text{CaCO}_3$  stimulated plant metabolism by enhancing photosynthesis and formation of plant pigments in ‘Crimson seedless’ grapevines and Morsy *et al.* (2008) also reported the same in pomegranates.

There are several calcium carbonate-based particle film products available for sunburn control in apples: Eclipse™ (Novazone Inc., Livermore, CA), Purshade® (Purfresh, Inc., Fremont, CA), Diffusion™ (Wilbur-Ellis Co., San Francisco, CA), Bud Mate (Agrichem Liquid Fertilizer Pty Ltd, Loganholme, QLD, Australia). In a comparative trial of Cocoon, Eclipse, Fruit Shield, Invelop, Raynox, Raynox Plus, Sun Guard, and Surround WP it was found that all materials increased the percentage of sunburn-free fruit, enabling packers to move more fruit into the highest grade and increasing the returns for individual growers (Hanrahan *et al.*, 2009). Furthermore, Purshade® significantly reduced sunburn when applied on three dates to ‘Granny Smith’ apples (<https://www.novasource.com/en/products/purshade>). These results are consistent with the results of the present study.

### Conclusions

In conclusion, under our experimental conditions, spraying trees with Parasol had a positive effect on the coloring of apples and also increased their firmness and the percentage of marketable fruit. The effectiveness of Parasol is likely related to several factors including a) the characteristics (origin, shape and size of particles) of the calcium carbonate, which optimizes the protective film due to its high opacifying; b) the ionic surfactant which ensures good coverage and creates a uniform film with remarkable stability; and c) the easy postharvest removal of the formed film.

### Literature Cited

- Ahmed, F.F., M.M. Shaaban, and A.M.K. Abd El-Aal. 2011. Protecting Crimson Seedless grapevines growing in hot climates from sunburn. *Res. J. Agri. Biol. Sci.* 7: 135-141.
- Alvarez, H.L., C.M. Di Bella, G.M. Colavita, P. Oricchio, and J. Strachnoy. 2015. Comparative effects of kaolin and calcium carbonate on apple fruit surface temperature and leaf net  $\text{CO}_2$  assimilation. *J. App. Hort.* 17: 176-180.
- Amarante, C.V.T., C.A. Steffens, and L.C. Argenta. 2011. Yield and fruit quality of ‘Gala’ and ‘Fuji’

- apple trees protected by white anti-hail net. *Sci. Hort.* 129: 79–85.
- Amerine, M.A. and C.S. Ough. 1980. *Methods for analysis of musts and wines*. Wiley, New York.
- Evans, R.G. 2004. Energy balance of apples under evaporative cooling. *Transact. Am. Soc. Eng.* 47: 1029–1037.
- Evans, R.G. and T.W. van der Gulik. 2011. Irrigation for microclimate control. p. 1015–1036. In: L. Stetson and A. Dedrick (eds.) *Irrigation*. 6th ed. The Irrigation Association, Falls Church, VA.
- Felicetti, D.A. and L.E. Schrader. 2008. Photooxidative sunburn of apples: Characterization of a third type of apple sunburn. *Int. J. Fruit Sci.* 8:160–172.
- Felicetti, D.A. and L.E. Schrader. 2009. Changes in pigment concentrations associated with sunburn browning of five apple cultivars. II. Phenolics. *Plant Sci.* 8:160–172.
- Glenn, D.M., E. Prado, A. Erez, J.R. McFerson, and G.J. Puterka. 2002. A reflective processed kaolin particle film affects fruit temperature, radiation reflection and solar injury in apple. *J. Amer. Soc. Hort. Sci.* 127: 188–193.
- Glenn, D.M. 2009. Particle film mechanisms of action that reduce the effect of environmental stress in ‘Empire’ apple. *J. Amer. Soc. Hort. Sci.* 134: 314–321.
- Hanrahan, I., T.R. Schmidt and J. McFerson. 2009. Comparison of commercial sunburn protection products. *HortScience* 44: 1084–1085.
- Iamsb, K., J. Sekozawa, S. Sugaya, H. Gemma, and J. Kamuro. 2009. Alleviating sunburn injury in apple fruit using natural and fertilizer forms of S-abscisic acid and its underlying mechanism. *J. Food Agric. Environ.* 7:446–452.
- Lal, N., and N. Sahu. 2017. Management strategies of sun burn in fruit crops – A review. *Int. J. Curr. Microbiol. App. Sci.* 6: 1126–1138.
- Lolicato, S. 2011. *Sunburn protection for fruit – a practical manual for orchardists in northern Victoria*. Dept. Environ. and Primary Ind., Victoria, Australia.
- McGuire, R. 1992. Reporting of objective color measurements. *HortScience* 27:1254–1255.
- Morsy, M., A.M.K. Abd El-Aal, and H.A. Abd El-Aal. 2008. Attempts to find the best preharvest treatment required for obtaining the optimum marketable fruits and its effect on storage life of ‘Manfalouty’ pomegranates. I. Evaluating of some soil and foliar treatments on splitting, sunburn, yield and fruit quality. *Minia J. Agric. Res. Develop.* 28: 263–283.
- Racsko, J. and L.E. Schrader. 2012. Sunburn of apple fruit: historical background, recent advances and future perspectives. *Critic. Rev. Plant Sci.* 31: 455–504.
- Reig, G., P. Jentsch, and D. Rosenberger. 2016. Effects of sunburn treatments on ‘Honeycrisp’ in the Hudson Valley in 2015. *New York Fruit Quarterly* 24(3): 5–10.
- Schrader, L., J. Zhang, and J. Sun. 2003. Environmental stresses that cause sunburn of apple. *Acta Hort.* 618: 397–405.
- Schmidt, T. 2018. Apple sunburn. 24 December 2019. <<http://treefruit.wsu.edu/article/apple-sunburn-101/>>.
- Schupp, J., E. Fallani, and Ik-Jo Chun. 2002. Effect of particle film on fruit sunburn, maturity and quality of ‘Fuji’ and ‘Honeycrisp’ apples. *Hort Technology*. 12: 87–90.  
<https://www.novasource.com/en/products/purshade>.